Inverse and Implicit Functions in Domain Theory

Abbas Edalat, Dirk Pattinson
Department of Computing
Imperial College, London UK
ae@doc.iac.ac.uk, dpattin@doc.ic.ac.uk

Abstract

We construct a domain-theoretic calculus for differentiable functions, which includes addition, subtraction and composition. Using the domain for differential functions, we develop a domain-theoretic version of the inverse function theorem and implicit function theorem, in which, subject to the usual conditions, the inverse function and the implicit function and their derivatives are obtained as fixed points of Scott continuous functionals and are approximated by step functions. This means that from an increasing sequence of step functions converging to a function and its derivative in the domain of differentiable functions we can effectively obtain an increasing sequence of step functions converging in this domain to the inverse function and its derivative, and also effectively obtain an increasing sequence of polynomial step functions whose lower and upper bounds converge in the C^1 norm to the inverse function. A similar result holds for implicit functions. Combined with the domain-theoretic model for computational geometry, this provides a robust technique for construction of curves and surfaces in geometric modelling and CAD.

1. Introduction

The earliest systematic method of computing increasingly better lower and upper bounds to π is credited to Archimedes, who obtained his approximation by circumscribing and inscribing the circle with regular polygons having 96 sides. In the 15th century, Jamshid Kashani, the Iranian mathematician, further developed this recursive method by using regular polygons with 805306368 sides, obtaining 16 decimal digits of π with a correct estimation of the error [1]. In today's mathematical language, we say that the two sequences of refining n-sided regular polygons circumscribing and inscribing the unit circle for $n = 3, 4, 5, \ldots$, as in Figure 1, converge in the C^1 norm respectively from outside and inside to the circle, a 1-dimensional closed manifold given by the implicit curve



Figure 1. Approximations of the unit circle

$x^2 + y^2 - 1 = 0$. This well-known and ancient technique describes the essence of our work in a simple and accessible way.

The present paper aims to develop a systematic and recursive method of approximating an implicitly given surface with two sequences of piecewise linear or piecewise polynomial surfaces, which converge locally in the C^1 norm from inside and outside to the implicit surface. A main application of this method is in geometric modelling and computer aided design, where curves and surfaces are usually defined implicitly [2]. Currently, there are no robust methods to approximate an implicit surface and the most reliable technique provided by interval analysis [12] is only able to approximate the implicit surface without approximating its derivative. The paper thus presents a framework for a robust CAD system, where implicitly given surfaces can be effectively obtained up to the C^1 precision required by the user.

In more mathematical language, we will use domain theory to provide a recursion-theoretic account of the inverse function and the implicit function theorems, which are the main fundamental tools in multi-variable differential calculus and the theory of manifolds.

In [7], a domain-theoretic framework for differential calculus of one variable was developed which in particular provides an effectively given domain for Lipschitz or differentiable functions. Later on, domain-theoretic techniques for solving initial value problems were obtained in [5, 9], which enable us to approximate the unique solution of an initial value problem given by a Lipschitz vector field up

to the precision required by the user. In [8], the domain-theoretic model was extended to multi-variable differential calculus, resulting in the notion of a domain-theoretic derivative, which for Lipschitz functions gives the smallest hyper-rectangle containing the Clark gradient [3], as well as a domain for multi-variable differentiable functions. In this paper, we will illustrate the first main application of the domain of multi-variable differentiable functions, by using it to obtain the inverse and implicit functions of classical analysis as the fixed point of a functional that we construct on the domain of differentiable functions. We first solve the inverse function problem by obtaining the inverse of a function near the identity map as a fixed point. Then, based on the domain-theoretic derivative, we introduce the notion of a mean differential and use it to solve the problem of inverse function in general. Finally the inverse function theorem is employed to obtain a domain-theoretic version of the implicit function theorem.

The domain-theoretic inverse and implicit function theorems have, in summary, the following distinguished features compared to other approaches:

(i) The inverse and implicit functions are obtained as fixed points of Scott continuous functionals.

(ii) From two increasing sequences of linear step functions converging to the function and its derivative, we can effectively obtain two increasing sequences of linear and polynomial step functions converging to the inverse of the function and its derivative. Similarly for the implicit function.

(iii) From two increasing sequences of linear step functions converging to the function and its derivative, we can effectively obtain an increasing sequences of polynomial step functions converging in the C^1 norm to the inverse of the function. Similarly, for the implicit function.

1.1 Related work

We have already mentioned the work on interval analysis regarding implicit surfaces; it gives approximations to the surface by voxel sets but there is no approximation of the derivative of the surface [12]. We here state the classical theorem on inverse functions for Lipschitz maps of Euclidean spaces, which has an elementary but non-constructive proof; see [10, Theorem 3.1].

For $x \in \mathbb{R}^n$ and the l_2 norm $\|\cdot\|_2$, we write $B_r = \{y \in \mathbb{R}^n \mid \|y\|_2 < r\}$ for the open ball around the origin with radius r and denote the closed ball by $\overline{B}_r\{y \in \mathbb{R}^n \mid \|y\|_2 \leq r\}$.

Theorem 1.1 Inverse Theorem for Lipschitz maps *[11] Let $\overline{B}_r$ be a closed ball containing the origin in $\mathbb{R}^n$ and let $f : \overline{B}r \to \mathbb{R}^n$ with $f(0) = 0$, so that for some invertible linear map $L : \mathbb{R}^n \to \mathbb{R}^n$ and some $\rho < 1$*

$$\|L^{-1}f(x_2) - L^{-1}f(x_1) - (x_2 - x_1)\|_2 \leq \rho\|x_2 - x_1\|_2$$

holds for all $x_1, x_2 \in \overline{B}_r$. Then, for all $x_1, x_2 \in \overline{B}_r$

$$\frac{1-\rho}{\|L^{-1}\|_2}\|x_2 - x_1\|_2 \leq \|f(x_2) - f(x_1)\|_2$$

$$\leq \|L\|_2(1+\rho)\|x_2 - x_1\|_2,$$

i.e. f is injective on $\overline{B}r$. The restriction $f|_{B_r} : B_r \to \mathbb{R}^n$ of f to the open ball B_r is an open map so that $V := f[B_r]$ is open. Thus $f|_{B_r} : B_r \to \mathbb{R}^n$ is Lipschitz with Lipschitz inverse and V contains the open ball B_s where $s = \frac{\rho(1-\rho)}{\|L^{-1}\|}$.

Note the non-constructive nature of the proof, which depends on the existence of a linear map L satisfying the above inequalities. We will actually see in the domain-theoretic Inverse Function Theorem 6.1 that, using the mean differential of the domain-theoretic derivative, we are able to compute a linear map L with the above property. A more general, but still non-constructive result, for the existence of an inverse of a Lipschitz function follows by using the Clarke gradient [3, p. 253], which is non-elementary. In Bishop's framework of constructive analysis, a constructive proof of the existence of the inverse function (or the implicit function) for a C^1 function is obtained by approximation but no approximations to the derivative of the inverse in provided [4]. In none of these approaches, the inverse or the implicit function is obtained as a fixed point of a functional.

2. Preliminaries

We briefly recall the essential notions of the domain-theoretic framework for multi-variable calculus from [8]. We write $D^0 =: [0,1]^n \to \mathbf{I}\mathbb{R}$ and $D^0_{\mathsf{i}} = \mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R}$, where $\mathbf{I}\mathbb{R}$ is the domain of non-empty compact intervals of $\mathbb{R}$, with the additional element $\perp = \mathbb{R}$, partially ordered with reverse inclusion. We identify a real number $x \in \mathbb{R}$ with the maximal element $\{x\} \in \mathbf{I}\mathbb{R}$ and a classical function $f : [0,1]^n \to \mathbb{R}$ with the function $x \mapsto \{f(x)\} : [0,1]^n \to \mathbf{I}\mathbb{R}$. Denote the smash product of n copies of a dcpo A by A^n_s, and let $\mathbf{I}\mathbb{R}^{n\times m}$ denote the collections of all $n \times m$ matrices with entries in $\mathbf{I}\mathbb{R}$.

The Scott continuous function $f : [0,1]^n \to \mathbf{I}\mathbb{R}$ has *an interval Lipschitz constant* $b \in (\mathbf{I}\mathbb{R})^{1\times n}_s$ in $a \in (\mathbf{I}[0,1])^n$ if for all $x, y \in a^\circ$ we have: $b(x-y) \sqsubseteq f(x) - f(y)$. The *single-step tie* $\delta(a,b) \subseteq D^0[0,1]$ of a with b is the collection of all functions in $D^0[0,1]$ which have an interval Lipschitz constant b in a. If $b \neq \perp$ then $\delta(a,b)$ consists only of functions which in the interior a° are classical Lipschitz functions. The well-defined and Scott continuous *primitive map* $\int : ([0,1]^n \to (\mathbf{I}\mathbb{R})^{1\times n}_s) \to (\mathcal{P}(D^0), \supseteq)$ is given by

$\int(\bigsqcup_{i\in I} a_i \searrow b_i) = \bigcap_{i\in I} \delta(a_i, b_i)$, where $\mathcal{P}$ is the power set functor. The *domain-theoretic derivative* of a continuous function $f : [0,1]^n \to \mathbf{I}\mathbb{R}$ is the well-defined and Scott continuous map $\frac{df}{dx} = \bigsqcup_{f\in\delta(a,b)} a \searrow b : [0,1]^n \to (\mathbf{I}\mathbb{R})^{1\times n}_s$. For a classical C^1 function f, we have $\frac{df}{dx} = f'$, where f' is the classical derivative of f, a notation which is used throughout the paper. For a classical Lipschitz map f, the domain-theoretic derivative $\frac{df}{dx}(x_0)$ at x_0 is is the smallest hyper-rectangle containing the Clarke gradient at x_0,which is a non-empty compact and convex set. The domain for Lipschitz functions is the continuous Scott subdomain

$$D^1([0,1[^n \to \mathbf{I}\mathbb{R}) \subset D^0([0,1]^n \to \mathbf{I}\mathbb{R}) \times D^0([0,1]^n \to \mathbf{I}\mathbb{R}^n_s)$$

of *consistent pairs* defined by $(f, g) \in D^1$ iff $\uparrow f \cap \int g \neq \emptyset$, which is equivalent to the existence of a classical continuous function $h : \mathrm{dom}(g) \to \mathbb{R}$ with $g \sqsubseteq \frac{dh}{dx}$. For $f \in D^0$, we have $(f, \frac{df}{dx}) \in D^1$, where $\frac{df}{dx}$ is, as always in this paper, the domain-theoretic derivative of f. In [8], it is shown that consistency is decidable on the rational step functions of $D^0([0,1]^n \to \mathbf{I}\mathbb{R}^n) \times D^0([0,1]^n \to \mathbf{I}\mathbb{R}^n_s)$ and thus D^1 can be given an effective structure. The framework can be extended in a straightforward way to functions of type $\mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R}$ and of type $\mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R}^m$.

Given $A \in \mathbf{I}\mathbb{R}^n$ and $B \in \mathbf{I}\mathbb{R}^m$, we write $C^1(A \to B)$ for the set of C^1 functions of type: $A \to B$. Similarly, we write $D^0(A \to \mathbf{I}B)$ for the continuous Scott domain $A \to \mathbf{I}B$, $D^0(\mathbf{I}A \to \mathbf{I}B)$ for the continuous Scott domain $\mathbf{I}A \to \mathbf{I}B$, and $D^1(A \to \mathbf{I}B)$ for the domain of differentiable functions. For $C \in \mathbf{I}\mathbb{R}^{n\times n}$, we also write $D^1(A \to \mathbf{I}B, A \to \mathbf{I}C)$ for consistent pairs in the subdomain $(A \to \mathbf{I}B) \times (A \to \mathbf{I}C)$. We write $D^0(A)$ for $D^0(A \to \mathbf{I}\mathbb{R})$ and $D^1(A)$ for $D^0(A \to \mathbf{I}\mathbb{R})$; also D^0 or D^1 stand $D^0[0,1]^n$ or $D^1[0,1]^n$, unless otherwise stated. For $f \in (A \to \mathbf{I}B)$ we define the *width* of f as $w(f) := \max_{1\leq i\leq m} sup_{x\in A} w(f_i(x))$ where $w(C)$ denotes the width of the interval C.

Recall the Scott continuous map $\mathcal{E} : D^0 \to D^0_{\mathsf{i}}$ with $\mathcal{E}(f) = \bigsqcap\{f(y) \mid y \in x\}$, which sends any map in D^0 to its maximal extension in D^0_{i}. For convenience, we sometimes write $\mathbf{I}f := \mathcal{E}(f)$. We also recall the notion of a piecewise polynomial step function [7]. Let $r \in \mathbf{I}[-1,1]^n$ and let $p, q : r \to \mathbb{R}$ be piecewise polynomial functions (i.e. each given by a finite number of polynomials) satisfying $p(x) \leq q(x)$ for $x \in r^\circ$. Note that p and q can be discontinuous in r°. The piecewise polynomial single-step functions $r \searrow [p,q] : [0,1]^n \to \mathbf{I}\mathbb{R}$ and $r \searrow_{\mathsf{i}} [p,q] : \mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R}$ are, respectively, given by

$$(r \searrow [p,q])(x) = \begin{cases} [p,q](x) & \text{if } x \in r^\circ \\ \bot & \text{otherwise.} \end{cases}$$

$$(r \searrow_{\mathsf{i}} [p,q])(x) = \begin{cases} \mathbf{I}[p,q](x) & \text{if } r \ll x \\ \bot & \text{otherwise.} \end{cases}$$

A piecewise polynomial step function is the lub $\bigsqcup_{i\in I} r_i \searrow [p_i, q_i]$ (or $\bigsqcup_{i\in I} r_i \searrow_{\mathsf{i}} [p_i, q_i]$) of a consistent finite set of piecewise polynomial single-step functions. If r_i, p_i and q_i are defined over rationals, then we say that the polynomial step function is rational. From now on we will be dealing with step functions of type $A \to \mathbf{I}B$ or $\mathbf{I}A \to \mathbf{I}B$ where $A, B \in \mathbf{I}\mathbb{R}^n$. A polynomial step function, say, of type $[0,1]^n \to \mathbf{I}\mathbb{R}^n$ will be given by a vector polynomial step function of type:

$$(r \searrow [p,q])(x) = \begin{cases} [p,q](x) & \text{if } x \in r^\circ \\ \bot & \text{otherwise.} \end{cases},$$

where $[p,q]$ is now a vector of n interval-valued functions given by polynomials $p_j \leq q_j$ for $j = 1, \cdots, n$. The composition $f \circ g$ of two piecewise linear step functions $f \in \mathbf{I}\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n$ and $g \in [0,1]^n \to \mathbf{I}\mathbb{R}^n$ is a piecewise linear step function of type $[0,1]^n \to \mathbf{I}\mathbb{R}^n$. The collection of rational piecewise linear (or polynomial) step functions gives a basis of $(D^0)^n_s$ and $(D^0_{\mathsf{i}})^n_s$.

The following gives an algorithm to evaluate $\mathcal{E}$ on a piecewise polynomial step function, which extends the algorithm in [7, Section 2]:

$$\mathcal{E}(\bigsqcup_{i\in I} r_i \searrow [p_i, q_i]) =$$
$$\bigsqcup\{(\sqcap_{j\in J} r_j) \searrow_{\mathsf{i}} [p_J, q_J] \mid J \subseteq I, \bigcup_{j\in J} r_j^\circ \text{ connected}\}, \quad (1)$$

where, for each $J \subset I$, the piecewise polynomials $p_J, q_J : (\sqcap_{j\in J} r_j) \to \mathbb{R}$ are defined for $x \in (\sqcap_{j\in J} r_j)$ by:

$$p_J(x) = \max\{p_j(x) \mid j \in J, x \in \mathrm{dom}(p_j)\}$$
$$q_J(x) = \min\{q_j(x) \mid j \in J, x \in \mathrm{dom}(q_j)\}.$$

Lemma 2.1 *If $\bigsqcup_{i\in I} r_i \searrow [p_i, q_i]$ is a piecewise linear step function, then so is $\mathcal{E}(\bigsqcup_{i\in I} r_i \searrow [p_i, q_i])$.*

Lemma 2.2 *If $u \in D^0$ and $u = \bigsqcup_{i\geq 0} s_i$, where s_i are piecewise linear step functions, then $\mathcal{E}(u) = \bigsqcup_{i\geq 0} \mathcal{E}(s_i)$ is the supremum of piecewise linear step functions.*

We will also need to use step functions made up of single-step functions of the form $c \searrow [p,q]$ and $c \searrow_{\mathsf{i}} [p,q]$ where $c \subseteq \mathbb{R}^n$ is a parallelogram. This class of step functions, say, of type $\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n$ is closed under pre-composition and post-composition with invertible linear maps:

$$(r \searrow [p,q]) \circ L = (L^{-1}r) \searrow [p,q],$$
$$L \circ (c \searrow [p,q]) = r \searrow [L \circ p, L \circ q],$$

and similarly:

$$(r \searrow_{\mathsf{i}} [p,q]) \circ L = (L^{-1}r) \searrow_{\mathsf{i}} [p,q]$$
$$L \circ (r \searrow_{\mathsf{i}} [p,q]) = r \searrow_{\mathsf{i}} [L \circ p, L \circ q].$$

We will use the maximum norm for vectors and matrices in $\mathbb{R}^n$, extended to interval valued vectors and matrices as follows. For $b \in \mathbf{I}\mathbb{R}^{m\times n}$, we define its maximum norm by $\|b\| = \max_{1\le i\le m}\sum_{j=1}^{n}\max\{|b_{ij}{}^-|, |b_{ij}^+|\}$, where $b_{ij} = [b_{ij}^-, b_{ij}^+]$. We write the maximum norm of a vector $s \in \mathbb{R}^n$ by $\|s\| := \max_{1\le i\le n}|s_i|$.

Definition 2.3 We say $f : \mathbf{I}A \to \mathbf{I}\mathbb{R}^m$ is *interval Lipschitz* in the open set $O \subset A$, if there exists $\ell \ge 0$ such that $w(f(x)) \le \ell w(x)$ for all $x \in \mathbf{I}[0,1]$ with $x \subset O$. We say that f is *interval contracting* if it is interval Lipschitz with an interval Lipschitz constant less than one.

Proposition 2.4 *If $A \in \mathbf{I}\mathbb{R}^n$, $f \in D^0(\mathbf{I}A \to \mathbf{I}\mathbb{R}^m)$ with $f \in \delta(a,b)$, $b \ne \bot$, then f has interval Lipschitz constant $n\|b\|$ in a°.*

Proof For any $x \in \mathbf{I}A$ with $a \ll x$, we have $b(x-x) \sqsubseteq f(x) - f(x)$. Since, for $y = [\underline{y}, \overline{y}] \in \mathbf{I}\mathbb{R}$, $y - y = [-(\overline{y} - \underline{y}), \overline{y} - \underline{y}]$ we obtain $w(y-y) = 2w(y)$. Now

$$\begin{aligned} 2nbw(x) &= \sum_{i=1}^{n} w([-bw(x), bw(x)]) \\ &\ge \sum_{i=1}^{n} w(b_i(x_i - x_i)) \\ &= w(b(x-x)) \\ &\ge w(f(x) - f(x)) = 2w(f(x)) \end{aligned}$$

which completes the proof.

Proposition 2.5 (i) *For $h \in D^0([0,1]^n \to \mathbf{I}\mathbb{R})$ and $g \in D^0([0,1]^n \to \mathbf{I}\mathbb{R}_s^n)$ we have $h \in \int g$ iff, for all $x, y \in [0,1]^n$ we have $\mathbf{I}g(x \sqcap y)(x-y) \sqsubseteq h(x) - h(y)$.*

(ii) *For $h \in D^0(\mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R})$ and $g \in D^0(\mathbf{I}[0,1]^n \to \mathbf{I}\mathbb{R}_s^n)$ we have $h \in \int g$ iff, for all $x, y \in \mathbf{I}[0,1]^n$ we have $g(x \sqcap y)(x-y) \sqsubseteq h(x) - h(y)$.*

Proof Similar to [7, Proposition 7.7].

Corollary 2.6 *Let $h \in D^0([0,1]^n \to \mathbf{I}\mathbb{R})$ and $g \in D^0([0,1]^n \to \mathbf{I}\mathbb{R}_s^n)$. Then $h \in \int g$ iff $\mathbf{I}h \in \int \mathbf{I}g$.*

Proof The "only if" part is trivial. For the "if part", let $X, Y \in \mathbf{I}[0,1]^n$, For $x \in X$ and $y \in Y$ we have by assumption:

$$\mathbf{I}g(x \sqcap y)(x-y) \sqsubseteq \mathbf{I}h(x) - \mathbf{I}h(y).$$

By taking glb over $x \in X$, we get:

$$\mathbf{I}g(X \sqcap y)(X-y) \sqsubseteq \mathbf{I}h(X) - \mathbf{I}h(y).$$

The result follows by taking glb over $y \in Y$. □

3. Domain-theoretic calculus of functions

We note that $f \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^m)$ iff $f_i \in D^1([0,1]^n \to \mathbf{I}\mathbb{R})$ for $i = 1, \cdots, n$, where $f = \begin{pmatrix} f_1 \\ \vdots \\ f_m \end{pmatrix}$.

For ease of presentation, we take the following notation: A vector $(x_1, \ldots, x_n) \in \mathbb{R}^n$ is denoted by $(x_i)_i$ a shorthand for $(x_i)_{1\le i\le n}$, i.e. a repeated index always runs from 1 to n,where n is the dimension of the Euclidean space $\mathbb{R}^n$, and describes the components of a vector, and similarly for a vector function. Thus, $f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$ is represented as $(f_0, (f_i)_i)$ where f_0 is the function part of f and $(f_i)_i$ its derivative part. This notation is also used for $n \times n$ matrices, i.e. $(a_{ij})_{ij}$ denotes an $n \times n$ matrix with ij entry a_{ij}. Thus, a vector function $f \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$ is denoted by $f = ((f_{i0})_i, (f_{ij})_{ij})$, where f_{i0} is the ith function part component of the f and $(f_{ij})_j$ for $j = 1, \ldots, n$ denotes the n derivative components of f_{i0}. The following statement is a version of the classical chain rule in the domain D^1.

Theorem 3.1 *Suppose $(f_0, (f_i)_i)) \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$ and $(g_{i0}, g_{i1}, \cdots, g_{im}) \in D^1([0,1]^m \to \mathbf{I}\mathbb{R})$ for $i = 1, \cdots n$. If*

$$h_0 = \mathbf{I}f_0((g_{i0})_i) \qquad h_j = (\sum_{k=1}^{n} \mathbf{I}f_k((g_{i0})_i) \cdot g_{kj}$$

for $1 \le j \le m$, then $(h_0, (h_j)_j) \in D^1([0,1]^m \to \mathbf{I}\mathbb{R})$.

Proof Let $h_i \in \uparrow g_{i0} \cap \int g_{i1}$, $k \in \uparrow f_0 \cap \int (f_1, \cdots, f_n)$ and put $t = k(h_1, \cdots, h_n)$. By Corollary 2.6, we have: $\mathbf{I}k \in \int(\mathbf{I}f_1, \cdots \mathbf{I}f_n)$. We show, by using Proposition2.5(i) that $t \in \int \sum_{i=1}^{n} f_i(g_{10}, \cdots, g_{n0}) \cdot g_{i1}$. For elements $x, y \in [0,1]$ we have:

$$\begin{aligned} & t(x) - t(y) \\ =& k(h_j(x)_j) - k(h_j(y)_j) \\ \sqsupseteq & \sum_{i=1}^{n} f_i((h_j(x) \sqcap h_j(y))j) \cdot (h_i(x) - h_i(y)) \\ \sqsupseteq & \sum_{i=1}^{n} f_i((h_j(x) \sqcap h_j(y))_j) \cdot g_{i1}(x \sqcap y)(x-y) \\ \sqsupseteq & \sum_{i=1}^{n} f_i((h_j(x \sqcap y))_j)) \cdot g_{i1}(x \sqcap y)(x-y) \\ \sqsupseteq & \sum_{i=1}^{n} f_i(g_{j0}(x \sqcap y)_j) \cdot g_{i1}(x \sqcap y)(x-y), \end{aligned}$$

which completes the proof.

We define the following arithmetic operations on $D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$.

Composition $(\cdot \circ \cdot)$:

$D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n) \times D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n) \to D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$

where, $f \circ g = ((h_{i0})_i, (h_{ij})_{ij})$ with

$$h_{i0} = \mathbf{I}f_{i0}(g_{m0})_m \qquad h_{ij} = \sum_{k=1}^{n} \mathbf{I}f_{ik}((g_{m0})_m) \cdot g_{kj}$$

for $1 \leq i, j \leq n$.

Addition $(\cdot + \cdot)$:

$D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^m) \times D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^m) \to D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^m)$

where$(f+g)_j = (f_{j0} + g_{j0}, f_{j1} + g_{j1}, \cdots f_{jn} + g_{jn})$.

Negation $(-)$:

$$D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n) \to D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n)$$

where $(-f)_j = (-f_{j0}, -f_{j1}, \ldots, -f_{jn})$.

Multiplication $(\cdot)$:

$$D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}) \times D^1(\mathbb{R} \to \mathbf{I}\mathbb{R}) \to D^1(\mathbb{R} \to \mathbf{I}\mathbb{R})$$

where $f \cdot g = (f_0 \cdot g_0, (f_i)_i \cdot g_0 + (g_i)_i \cdot f_0)$.

Inversion $(\frac{1}{\cdot})$:

$$D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}) \to D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$$

where $\frac{1}{f} = (\frac{1}{f_0}, \frac{-f_1}{f_0^2}, \ldots, \frac{-f_n}{f_0^2})$ and, as usual, for intervals a and b, we put $a/b = \perp$ if $0 \in b$.

Corollary 3.2 *The arithmetic operations are well-defined:*

(i) *If $f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n)$ and $g \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$, then $f \circ g \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$.*

(ii) *If $f \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$ and $g \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$, then $f + g \in D^1([0,1]^n \to \mathbf{I}\mathbb{R}^n)$.*

(iii) *If $f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n)$, then $-f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n)$.*

(iv) *If $f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$ and $g \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$, then $f \cdot g \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$.*

(v) *If $f \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$, then $\frac{1}{f} \in D^1(\mathbb{R}^n \to \mathbf{I}\mathbb{R})$.*

Moreover these operations are Scott continuous.

Lemma 3.3 Chain Rule. *For any two functions $f, g \in D^0(\mathbb{R}^n \to \mathbf{I}\mathbb{R}^n)$:*

$$\frac{d(f \circ g)}{dx} \sqsupseteq (\frac{df}{dx} \circ g) \cdot \frac{dg}{dx}.$$

Proof Fix $x_0 \in \mathbb{R}^n$. Let $b_1 \ll \frac{dg}{dx}(x_0)$ and $b_2 \ll (\frac{df}{dx}(g(x_0))$. Then for some $a_1, a_2 \in \mathbf{I}\mathbb{R}^n$ with $x_0 \in a_1$ and $g(x_0) \in a_2$ we have: $f \in \delta(a_1, b_1)$ and $g \in \delta(a_2, b_2)$. Since g is Scott continuous $g^{-1}(\Uparrow a_2)$ is open in $\mathbb{R}^n$. Let $a \in \mathbf{I}\mathbb{R}^n$, with $a \subseteq a_1 \cap g^{-1}(\Uparrow a_2)$ and $g(x_0) \in a^\circ$. Then, for all $x, y \in a^\circ$:

$$f(g(x)) - f(g(y)) \sqsupseteq b_1(g(x) - g(y)) \sqsupseteq b_1 b_2(x - y),$$

and it follows that $f \circ g \in \delta(a, b_1 b_2)$ and thus $\frac{d(f \circ g)}{dx}(x_0) \sqsupseteq b_1 b_2$.

Note that equality may fail in the chain rule. For example let $f : \mathbb{R} \to \mathbb{R}$ be the absolute value function $x \mapsto |x|$ and $g : \mathbb{R} \to \mathbb{R}$ defined by $x \mapsto x$ if $x \leq 0$ and 0 if $x > 0$. Then $\frac{df \circ g}{dx}(0) = [-1, 0]$ whereas $\frac{df}{dx}(g(0)) \cdot \frac{dg}{dx}(0) = \frac{df}{dx}(0) \cdot \frac{dg}{dx}(0) = [-1, 1][0, 1] = [-1, 1]$.

Lemma 3.4 *A map $f : [-a, a]^n \to \mathbb{R}^n$ has Lipschitz constant nc with respect to the max norm on $\mathbb{R}^n$ iff $f \in (\int [-a, a]^n \searrow [-c, c]^n)^n$.*

Proof Suppose f has Lipschitz constant nc. Then for each $i = 1, \cdots, n$ we have: $|f_i(x) - f_i(y)| \leq nc|x - y|$. If m is such that $|x_m - y_m| = |x - y|$, then we obtain:

$$f_i(x) - f_i(y) \sqsupseteq [-nc, nc](x_m - y_m) \sqsupseteq \sum_{j=1}^{n} [-c, c](x_j - y_j),$$

are required. On the other hand, if $f \in (\int [-a, a]^n \searrow [-c, c]^n)^n$, then for each $i = 1, \cdots, n$ we have:

$$f_i(x) - f_i(y) \sqsupseteq \sum_{j=1}^{n} [-c, c](x_j - y_j),$$

and thus:

$$|f_i(x) - f_i(y)| \leq \sum_{j=1}^{n} c|x_j - y_j| \leq nc|x - y|,$$

which yields: $|f(x) - f(y) \leq nc|x - y|$.

Lemma 3.5 *Let $(g_0, g_1, \cdots, g_n) \in D^1([0,1]^n \to \mathbf{I}\mathbb{R})$ with g_0 real valued. If $g_i(x)$ is a real number for all $i = 1, \cdots, n$ for some $x \in [0,1]^n$, then $g_0'(x)$ exists and $g_0'(x) = (g_1(x), \cdots, g_n(x))$.*

Proof By assumption, there exists $t \in \uparrow g_0 \cap \int(g_1, \cdots, g_n)$, but then $t = g_0$ as g_0 is maximal. It follows that $\frac{dg_0}{dx} \sqsupseteq (g_1, \cdots, g_n)$, which implies $\frac{dg_0}{dx}(x) = (g_1(x), \cdots, g_n(x)))$. By [7, Proposition 4.3(ii)], the result follows.

4. Inverse constructing functional

In many areas of mathematics a construction is first obtained for functions close to the identity map and then extended to more general maps. Here, we show first that the inverse of a function close to the identity function can be obtained as the fixed point of a functional that we introduce here. Later in the paper, we show how this functional can be used to obtain the inverse of a Lipschitz function.

For $a > 0$ and $0 < c \leq 1/2$, let $A = [-a, a]^n$ and $B = [-ca, ca]^n$, and consider the functional

$$T : D^1(A \to \mathbf{I}B) \times D^1(B \to \mathbf{I}B) \to D^1(B \to \mathbf{I}B)$$

defined by

$$T(f, g) = -\mathbf{I}f(I + g)$$

where $I : B \to \mathbf{I}B$ is the identity function $\lambda x.x$. We also put $T_f = T(f, \cdot)$. By Corollary 3.2 T is well-defined. Later, we construct the inverse function of f as the least fixpoint of T_f. The function part of T is given by the functional

$$\begin{array}{lccc} R: & D^0(A \to \mathbf{I}B) \times D^0(B \to \mathbf{I}B) & \to & D^0(B \to \mathbf{I}B) \\ & (f, g) & \mapsto & -\mathbf{I}f \circ (I + g) \end{array}$$

Let $R_f = R(f, \cdot)$. For the derivative part, we have the functional S of type

$$D^0(A \to \mathbf{I}B) \times (B \to \mathbf{I}B) \times (B \to \mathbf{I}\mathbb{R}^n) \to (B \to \mathbf{I}\mathbb{R}^n)$$

defined by

$$S(h, g_0, g_1) = -\mathbf{I}h \circ (I + g_0) \cdot (E + g_1)$$

where $E = \frac{dI}{dx}$ is the identity matrix. We also put $S_{(h,g_0)} = S(h, g_0, \cdot)$. Thus,

$$T(f, g) = (R((f_{i0})_i, (g_{i0})_i), S((f_{ij})_{ij}, (g_{i0})_i, (g_{ij})_{ij})$$

if $f = ((f_{i0})_i, (f_{ij})_{ij}) \in D^1(A \to \mathbf{I}B)$ and $g = ((g_{i0})_i, (g_{ij})_{ij}) \in D^1(B \to \mathbf{I}B)$.

Proposition 4.1 *Suppose $f : [-a, a]^n \to \mathbb{R}^n$. Then $R_f : D^0([-ca, ca]^n \to \mathbf{I}[-ca, ca]^n) \to D^0([-ca, ca]^n \to \mathbf{I}[-ca, ca]^n)$ is well defined. If moreover f has Lipschitz constant $c < 1/n$ for $n > 1$ ($c \leq 1/2$ for $n = 1$) with respect to the max norm and $f(0) = 0$, we have:*

(i) *The functional R_f has a unique fixed point $(g_{i0})_i$; it satisfies: $g_{i0}(0) = 0$ and $w(g_{i0}) = 0$ for $1 \leq i \leq n$, and is thus a classical C^0 function.*

(ii) *$I + f : [-a, a]^n \to \mathbb{R}^n$ has inverse $I + (g_{i0})_i : Im(I + f) \to [-a, a]^n$.*

Proof It is easily checked that R_f is well-defined using $c < 1/n$ for $n > 1$ ($c \leq 1/2$ for $n = 1$). For (i), let $(g_{i0})_i$ be the least fixed point of R_f, then we have:

$$g_{i0} = -\mathbf{I}f_i(\pi_k + g_k)_k \tag{2}$$

By Lemma 3.4, $f_i \in \int[-a, a]^n \searrow [-c, c]^n$ for each $i = 1, \cdots, n$, which implies $\mathbf{I}f_i \in \int[-a, a]^n \searrow [-c, c]^n$. Thus, by Proposition 2.5, we have from Equation 2:

$$\begin{array}{rcl} -g_i(0) & = & \mathbf{I}f_i(g_1(0), \cdots, g_n(0)) - \mathbf{I}f_i(0) \\ & \sqsupseteq & \sum_{i=1}^n [-c, c] g_i(0). \end{array}$$

Thus, $\|g_i(0)\| \leq c(\sum_{i=1}^n \|g_i(0)\|)$ for each $i = 1, \cdots, n$. Summing up for $i = 1, \cdots, n$, we obtain: $\sum_{i=1}^n \|g_i(0)\| \leq nc(\sum_{i=1}^n \|g_i(0)\|)$. Since $c < 1/n$ for $n > 1$, it follows that $g_i(0) = 0$ for each $i = 1, \cdots, n$. By Proposition 2.4, $\mathbf{I}f_i$ is contracting with contractivity factor c for each $i = 1, \cdots, n$. Let $x \in [-ca, ca]^n$, we have

$$\begin{aligned} w(g_i(x)) &= w(\mathbf{I}f_i((x_k + g_k(x)))_k) \\ &\leq c \max_{1 \leq i \leq n} w(x_i + g_i(x))) \\ &= c \max_{1 \leq i \leq n} w(g_i(x)). \end{aligned}$$

Summing for $i = 1, \cdots, n$, we obtain:

$$\max_{1 \leq i \leq n} w(g_i(x)) \leq \sum_{1 \leq i \leq n} w(g_i(x)) \leq nc \max_{1 \leq i \leq n} w(g_i(x))$$

Since $nc < 1$, it follows that $\max_{1 \leq i \leq n} w(g_i(x)) = 0$. It then follows that $w(g_i) = 0$. Thus, the least fixed point g is maximal and hence the unique fixed point.

(ii) We have $(I + f) \circ (I + g) = I + g + f \circ (I + g) = I$ and thus $I + g$ is a right inverse of $I + f$. Since $I + f$ is Lipschitz with $\|f(x) - f(y)\| \leq nc\|x - y\|$ it follows by [10, Theorem 3.1] that $I + f$ has an inverse. Thus, $I + g$, being a right inverse, is the inverse of $I + f$.

Now we can obtain a fixed point of $T_{(f, \frac{df}{dx})}$ and examine its properties.

Lemma 4.2 *The partial order $(D^0([0, 1] \to \mathbf{I}\mathbb{R}), \sqsupseteq)$ of Scott continuous functions ordered by reverse pointwise ordering is a dcpo with Scott continuous operations for addition $\cdot + \cdot : D^0([0, 1] \to \mathbf{I}\mathbb{R}) \times D^0([0, 1] \to \mathbf{I}\mathbb{R})$, multiplication $\cdot \times \cdot : D^0([0, 1] \to \mathbf{I}\mathbb{R}) \times D^0([0, 1] \to \mathbf{I}\mathbb{R})$ and negation $-\cdot : D^0([0, 1] \to \mathbf{I}\mathbb{R}) \to D^1([0, 1] \to \mathbf{I}\mathbb{R})$.*

Proof Note that any bounded complete dcpo is a dcpo with respect to its opposite order. It is routine to check the Scott continuity of addition and multiplication.

Proposition 4.3 *Suppose $f : [-a, a]^n \to \mathbb{R}^n$ has Lipschitz constant $c < 1/n$ for $n > 1$ ($c \leq 1/2$ for $n = 1$) and $f(0) = 0$.*

(i) *The functional*

$$T_{(f,\frac{df}{dx})} : D^1(X_0, X_1) \to D^1(X_0, X_1),$$

where $X_0 = [-ca, ca]^n \to [-ca, ca]^n$ *and* $X_1 = ([-ca, ca]^n \to \mathbf{I}[-d, d]^n)^n$*, with* $d = \frac{nc}{1-nc}$*, is well-defined and has a unique fixed point* $g = ((g_{i0})_i, (g_{ij})_{ij})$*, such that* $(g_{i0})_i$ *is the unique fixed point of* R_f *and* $(g_{ij})_{ij}$ *is the unique fixed point of* $S_{(\frac{df}{dx},(g_{i0})_i)} : X_1 \to X_1$*; moreover* $(g_{ij})_{ij} = \bigsqcap_{l\geq 0} S^l_{(\frac{df}{dx},(g_{i0})_i)}(\frac{d(g_{i0})_i}{dx}) \sqsubseteq \frac{d(g_{i0})_i}{dx}$.

(ii) *The fixed point* g *of* $T_{(f,\frac{df}{dx})}$ *will satisfy* $w((g_{ij}(x))_{ij}) \leq \frac{n(1+d)}{1-nc} w(\frac{df}{dx}(x_k + g_{k0}(x))_k)$ *for any* $x \in [-ca, ca]^n$ *and thus,* $w(g_{ij})_{ij} \leq \frac{n(1+d)}{1-nc} w(\frac{df}{dx})$.

(iii) *If* $w(\frac{df}{dx}((x_k + g_{k0}(x))_k) = 0$ *for some* $x \in [-ca, ca]^n$*, then* $w(g_{ij}(x)) = 0$*, for* $i, j = 1, \cdots, n$*, and* $(g_{i1}(x), \cdots, g_{in}(x)) = g'_{i0}(x)$ *for* $i = 1, \cdots, n$.

(iv) *If* $w(\frac{df}{dx}) = 0$*, then* $w(g_{i0}) = 0$ *for* $i = 1, \cdots n$*, and* $(g_{i1}, \cdots, g_{in}) = g'_{i0}$ *for* $i = 1, \cdots n$.

(iv) *If* $\frac{df}{dx}(0) = 0$ *then* $g_{ij}(0) = 0$ *for* $i, j = 1, \cdots, n$.

Proof (i) That $T_{(f,\frac{df}{dx})}$ is well-defined follows from $c < 1/n$. First, assume $g = ((g_{i0})_i, (g_{ij})_{ij})$, with $g_{ij} \neq \perp$ $(i, j = 1, \ldots, n)$, is any fixed point of the extended functional

$$T_{(f,\frac{df}{dx})} : D^1(X_0, X'_1) \to D^1(X_0, X'_1)$$

where $X'_1 = D^0([-ca, ca]^n \to \mathbf{I}\mathbb{R}^n)$. The fixed point equation gives us:

$$g_{i0} = -\mathbf{I}f_{i0}((\pi_k + g_{k0})_k) \tag{3}$$

$$g_{ij} = \sum_{m=1}^{n} -\mathbf{I}(\frac{df}{dx})_{im}((\pi_k + g_{k0})_k) \cdot (\lambda x.\delta_{mj} + g_{mj}). \tag{4}$$

Thus, $(g_{i0})_i$ is the unique fixed point of R_f. Moreover, by Lemma 3.4, we have $f \in (\int[-a, a]^n \searrow [-c, c]^n)^n$ and thus $\frac{df}{dx} \sqsupseteq [-a, a]^n \searrow [-c, c]^n$. Hence, we have for each $i = 1, \cdots, n$:

$$\|g_{ij}\| \leq c(\sum_{m=1}^{n} \|\delta_{mj} + g_{mj}\|) \leq c(1 + \sum_{m=1}^{n} \|g_{mj}\|).$$

Summing for $i = 1, \cdots, n$, we obtain:

$$\sum_{m=1}^{n} \|g_{mj}\| \leq \frac{nc}{1-nc}.$$

Therefore, $(g_{ij})_{ij} \sqsupseteq [-ca, ca]^n \searrow [-\frac{nc}{1-nc}, \frac{nc}{1-nc}]$. Hence, to prove the existence of the least fixed point $((g_{i0})_i, g_{\min})$ of $T_{(f,\frac{df}{dx})}$, it is sufficient to show that:

$$(\perp_0, \perp_1) \sqsubseteq T_{(f,\frac{df}{dx})}(\perp_0, \perp_1),$$

where $\perp_0 = [-ca, ca]^n \searrow [-ca, ca]^n$ and $\perp_1 = ([-ca, ca]^n \searrow [-\frac{nc}{1-nc}, \frac{nc}{1-nc}])^n$. We only need to check the information refinement for the derivative approximation. We put

$$(h_{ij})_{ij} = T_{(f,\frac{df}{dx})}(\perp_0, \perp_1).$$

Then, for $1 \leq i, j \leq n$,

$$\begin{aligned} h_{ij} &\sqsupseteq \sum_{m=1}^{n} [-c, c](1 + [\frac{-nc}{1-nc}, \frac{nc}{1-nc}]) \\ &= [\frac{-nc(1-nc+nc)}{1-nc}, \frac{nc(1-nc+nc)}{1-nc}] \\ &= [\frac{-nc}{1-nc}, \frac{nc}{1-nc}] = [-d, d], \end{aligned}$$

as required. For any fixed point $((g_{i0})_i, (g_{ij})_{ij})$ we have $(g_{ij})_{ij} \sqsubseteq \frac{d(g_{i0})_i}{dx}$ since $((g_{i0})_i, (g_{ij})_{ij})$ is consistent and $(g_{i0})_i$ has zero width. Now for any $h \in X_1$ with $g_{\min} \sqsubseteq h \sqsubseteq \frac{d(g_{i0})_i}{dx}$, we have, by the Chain rule 3.3:

$$g_{\min} \sqsubseteq S_{(\frac{df}{dx},(g_{i0})_i)}(h) \sqsubseteq S_{(\frac{df}{dx},(g_{i0})_i)}(\frac{d(g_{i0})_i}{dx}) \sqsubseteq \frac{d(g_{i0})_i}{dx}.$$

It follows by Lemma 4.2 that, in the lattice of functions in X_1 with $\sqsupseteq$ as the ordering, $g_{\min}$ as the top element and $\frac{d(g_{i0})_i}{dx}$ as the least element, the Scott continuous function $S_{(f,(g_{i0})_i)} : X_1 \to X_1$ has a least fixed point

$$g_{\max} = \bigsqcap_{l\geq 0} S^l_{(\frac{df}{dx},(g_{i0})_i)}(\frac{d(g_{i0})_i}{dx}),$$

which is the greatest fixed point of $S_{(f,(g_{i0})_i)} : X_1 \to X_1$ and thus induces the greatest fixed point $((g_{i0})_i, g_{\max})$ of $T_{(f,\frac{df}{dx})}$. It remains to show the uniqueness of the fixed point of $S_{(f,(g_{i0})_i)} : X_1 \to X_1$.

The fixed point equation (4) evaluated at $x \in [-ca, ca]^n$ yields:

$$g_{ij}(x) = \sum_{m=1}^{n} -\mathbf{I}(\frac{df}{dx})_{im}(x_k + g_{k0}(x))_k \cdot (\delta_{mj} + g_{mj}(x)),$$

or equivalently,

$$\begin{aligned} [g^-_{ij}(x), g^+_{ij}(x)] = \sum_{m=1}^{n} ([c_{im}(x), d_{im}(x)] \cdot \\ [\delta_{mj} + g^-_{mj}(x), \delta_{mj} + g^+_{mj}(x)]), \end{aligned} \tag{5}$$

where

$$[c_{im}(x), d_{im}(x)] := -\mathbf{I}(\frac{df}{dx})_{im}(x_k + g_{k0}(x))_k),$$

for $1 \leq i, m \leq n$. For any fixed $x \in [-ca, ca]^n$, consider the system of $2n^2$ equations represented by:

$$[y_{ij}, z_{ij}] = \sum_{m=1}^{n} [c_{im}(x), d_{im}(x)] \cdot [\delta_{mj} + y_{ij}, \delta_{mj} + z_{ij}], \tag{6}$$

for the unknown values y_{ij} and z_{ij} $(i, j = 1, \cdots, n)$. This is a linear system, which can be written in the form:

$$A(x)u = b(x) \tag{7}$$

where $A(x) \in \mathbb{R}^{2n^2 \times 2n^2}$ and $u, b(x) \in \mathbb{R}^{2n^2}$, with $u_{0ij} = y_{ij}$ and $u_{1ij} = z_{ij}$, in which the subscripts of u are numbers written in base n, i.e., $kij = kn^2 + in + j$. We already know that this linear system has at least one solution since $S_{(f,(g_{i0})_i)} : X_1 \to X_1$ has at least one fixed point. Suppose, for a contradiction, that the fixed point is not unique. Then $(g_{\min})_{i_0 j_0}(x_0) \neq (g_{\max})_{i_0 j_0}(x_0)$ for some $x_0 \in [-ca, ca]^n$ and some i_0, j_0 with $1 \leq i_0, j_0 \leq n$. And thus, for $x = x_0$, the linear system (7) will have infinitely many solutions of the form:

$$u_{0ij}(k) = (g_{\min})^-_{ij}(x_0) + \sum_{t=1}^{p} k_t \theta^t_{0ij}$$
$$u_{1ij} = (g_{\min})^+_{ij}(x_0) + \sum_{t=1}^{p} k_t \theta^t_{1ij},$$

where $\theta^t \in \mathbb{R}^{2n^2}$ $(t = 1, \cdots p)$ is a basis for the null-set of A and $k \in \mathbb{R}^p$. But $(g_{\min})_{ij}(x_0) \supseteq (g_{\max})_{ij}(x_0)$ and for some $k \in \mathbb{R}^p$, we have:

$$u_{0i_0j_0}(k) = (g_{\max})^-_{i_0j_0}(x_0), \qquad u_{1i_0j_0} = (g_{\max})^+_{i_0j_0}(x_0).$$

Now, using this value of k, we define $h : [-ca, ca]^n \to \mathbf{I}[-d, d]^{n \times n}$ by

$$h_{ij}(x) = \begin{cases} (g_{\min})_{i_0j_0}(x_0) - \sum_{t=1}^{p} k_t [\theta^t_{0i_0j_0}, \theta^t_{1i_0j_0}] \\ \quad \text{if } i = i_0, j = j_0, x = x_0 \\ (g_{\min})_{ij}(x) \quad \text{otherwise.} \end{cases}$$

Then, h is Scott continuous and is a fixed point of $S_{(f,(g_{i0})_{1 \leq i \leq n})} : X_1 \to X_1$ with $h \sqsubseteq g_{\min}$ and $h \neq g_{\min}$, which is the required contradiction.

(ii) We have,

$$\begin{aligned} & w(g_{ij}(x)) \\ =& w(\sum_{m=1}^{n} -\mathbf{I}(\frac{df}{dx})_{im}(x_k + g_{k0}(x))_k) \cdot (\delta_{mj} + g_{mj}(x)) \\ \leq& \sum_{m=1}^{n} w(\frac{df}{dx})_{im}((x_k + g_{k0}(x))_k)(1 + d) + cw(g_{mj}(x)) \\ \leq& n(1 + d)w(\frac{df}{dx})_i((x_k + g_{k0}(x))_k) + c\sum_{m=1}^{n} w(g_m(x)) \\ \leq& n(1 + d)w(\frac{df}{dx}((x_k + g_{k0}(x))_k)) + ncw(g(x)). \end{aligned}$$

Thus, $w(g(x)) \leq n(1 + d)w(\frac{df}{dx}((x_k + g_{k0}(x))_k)) + ncw(g(x))$, which gives: $w(g(x)) \leq \frac{n(1+d)}{1-nc} w(\frac{df}{dx}((x_k + g_{k0}(x))_k))$.

(iii) From (ii), $w(g_{ij}) = 0$ for $i, j = 1, \cdots, n$. Lemma 3.5, gives $(g_{i1}(x), \cdots, g_{in}(x)) = g'_{i0}(x)$ for $i = 1, \cdots, n$.

(iv) Follows from (iii).

(iv) Since $\mathbf{I}\frac{df}{dx}(0) = 0$, from Equation (4) we get: $g_{ij}(0) = 0$ for $i, j = 1, \cdots, n$.

Corollary 4.4 *Suppose $f \in C^1([-a, a]^n \to [-ca, ca]^n)$ with $\|f'(x)\| < \min\{\frac{1}{2}, \frac{1}{n}\}$ for all $x \in [-a, a]^n$ and assume $f(0) = f'(0) = 0$. Then, there is a unique fixed point $((g_{i0})_i), (g_{ij})_{ij})$ of $T_{(f,f')}$. It will satisfy $w(g_{i0}) = w(g_{ij}) = 0$ and $g_{i0}(0) = g_{ij}(0) = 0$ for $1 \leq i \leq n, 0 \leq j \leq n$, and $(g_{ij})_{ij} = (g_{i0})'_i$.*

5 The mean differential

In the previous section, we have constructed a local inverse for functions whose domain-theoretic derivative was close to zero in norm. In this section, we establish the necessary apparatus to extend this result to functions with arbitrary, but non-zero domain theoretic derivative. This is established by multiplying the function with a suitable matrix, the inverse of the mean differential, introduced below.

For $b \in \mathbf{I}\mathbb{R}^{m \times n}$, let $M \in \mathbb{R}^{m \times n}$ be given by $M_{ij} = ({b_{ij}}^- + {b_{ij}}^+)/2$, which we call the *mid-point* matrix of b. We say $b \in \mathbf{I}\mathbb{R}^{m \times n}$ is *zero centred* if for each i, j $(1 \leq i \leq m$ and $1 \leq j \leq n)$, $b_{ij} = [-r_{ij}, r_{ij}]$ for some $r_{ij} \geq 0$. If $b = (b_{ij}) \in \mathbf{I}\mathbb{R}^{m \times n}$ and $L = (L_{ij})_{ij} \in \mathbb{R}^{m \times n}$, we write $L \in b$ if $L_{ij} \in b_{ij}$ for all $1 \leq i \leq m, 1 \leq j \leq n$.

Lemma 5.1 *Suppose $b \in \mathbf{I}\mathbb{R}^{n \times n}$ and the $n \times n$ matrix $L \in b$ is invertible such that $L^{-1}b - I$ is zero-centred. Then $L = M$ the mid-point matrix of b.*

Proof Writing down what it means to say that $L^{-1}b - I$ is zero-centered, we obtain $L^{-1}M - I = 0$, from which the result follows immediately.

Proposition 5.2 *If $0 \notin \det b$ then for any $L \in b$, we have $\|M^{-1}b - I\| \leq \|L^{-1}b - I\|$, where M is the mid-point matrix of b.*

Proof Let the elements of L be given by l_{ij} and let (ℓ_{ij}) be the matrix of the cofactors of L so that $(L^{-1})_{ij} = \ell_{ji}/\det L$. The ij entry of $L^{-1}b - I$ is the interval:

$$\left[\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{-\sigma(\ell_{mi})}}{\det L} - \delta_{ij}, \frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right],$$

where $\sigma(r)$ is the sign of the real number r. Note that for any $L \in b$, the zero matrix is contained in $L^{-1}b - I$ and thus zero is contained in each of the above intervals for $i, j = 1, \cdots, n$. Recall that both $\det L$ and ℓ_{ij} (for each pair $i, j = 1, \cdots, n$) are linear in l_{pq} for $p, q = 1, \cdots n$, and thus, for each pair $p, q = 1, \cdots, n$, the two maps

$$l_{pq} \mapsto \left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{-\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right|$$

$$l_{pq} \mapsto \left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right|,$$

are linear fractional transformations, one increasing and one decreasing for $l_{pq} \in b_{pq}$. It follows that the minimum of

$$\max\left(\left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{-\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right|, \left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right|\right)$$

is at the point when the two linear fractional transformations coincide in value. We conclude that the minimum value of $\|L^{-1}b - I\|$ is reached when

$$\left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{-\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right| = \left|\frac{\sum_{m=1}^n \ell_{mi} b_{mj}^{\sigma(\ell_{mi})}}{\det L} - \delta_{ij}\right|$$

for all $ij = 1, \cdots n$, i.e., when $L^{-1}b - I$ is zero-centered. By Lemma 5.1 it follows that this minimum is achieved for $L = M$. □

Lemma 5.3 *If there is an invertible $L \in b$ such that $\|L^{-1}b - I\| < 1$, then $0 \notin \det b$.*

Proof Suppose such an L exists. Then we claim that for any $K \in b$ the n columns of $C = L^{-1}K$ are linearly independent. In fact suppose there exist $s_i \in \mathbb{R}$ $(j = 1, \cdots, n)$ with $\sum_{j=1}^n s_j C_{ij} = 0$ for all $i = 1, \cdots, n$. Let m be such that $|s_m| \geq |s_j|$ for all $j = 1, \cdots, n$. If $s_m \neq 0$, then we have:

$$\sum_{j=1}^n \frac{s_j}{s_m} C_{mj} = 0.$$

Thus,

$$1 = |C_{mm} - 1 + \sum_{j \neq m} \frac{s_j}{s_m} C_{mj} \leq |C_{mm} - 1| + \sum_{j \neq m} |C_{mj}| < 1,$$

a contradiction. It follows that C is invertible and thus so is K and the proof is complete. □

Corollary 5.4 *Let $0 \leq c < 1$. The following two conditions are equivalent.*

(i) *There exists an invertible $L \in b$ such that $\|L^{-1}b - I\| \leq c$.*

(ii) *The mid-point matrix M of b is invertible and $\|M^{-1}b - I\| \leq c$.*

Now consider a locally Lipschitz map $u : [-1, 1]^n \to \mathbb{R}^n$. Then the domain-theoretic derivative $\frac{du}{dx}(x_0)$ at a point $x_0 \in [-1, 1]$ is an $n \times n$ matrix of compact intervals. We can now introduce a simple notion for the domain-theoretic derivative:

Definition 5.5 The linear map represented by the mid-point matrix of $\frac{du}{dx}(x_0)$ is called the *mean differential* of u at x_0.

We will see in the next section that the mean differential is a main tool to develop the domain-theoretic version of the inverse function theorem for Lipschitz maps, which refines the classical result.

6. Inverse functions

We now prove the following theorem, which refines the classical result on inverse functions for Lipschitz functions 1.1.

Theorem 6.1 Inverse Function Theorem. *Let $u : [-1, 1]^n \to \mathbb{R}^n$ be locally Lipschitz with $u(0) = 0$. Suppose the mean differential at the point 0 (i.e., the linear map represented by the mid-point matrix of $\frac{du}{dx}(0)$), denoted by M, is invertible. Let $\|M^{-1}\frac{du}{dx}(0) - I\| < \min\{1/n, 1/2\}$. Then we have:*

(i) *The map u has a Lipschitz inverse in a neighbourhood of the origin.*

(ii) *If two increasing sequences of linear rational step functions, respectively of type $[-1, 1]^n \to \mathbf{I}\mathbb{R}^n$ and $[-1, 1]^n \to \mathbf{I}\mathbb{R}^{n^2}$, converging respectively to u and $\frac{du}{dx}$ are given, then we can effectively obtain an increasing sequence of piecewise linear step functions converging to the inverse of u.*

(iii) *If u is C^1 and two increasing sequences of linear rational step functions, respectively of type $[-1,1]^n \to \mathbf{I}\mathbb{R}^n$ and $[-1,1]^n \to \mathbf{I}\mathbb{R}^{n^2}$, converging respectively to u and u' are given, then we can also effectively obtain an increasing sequence of polynomial step functions converging to the derivative of the inverse of u.*

(iv) *With the assumptions in (iii), we can also effectively obtain an increasing sequence of polynomial step functions, whose lower and upper parts are continuous and piecewise polynomial, such that the two sequences of lower and upper parts converge in the C^1 norm, respectively from below and above, to the inverse of u.*

Proof (i) Suppose $N \in \frac{du}{dx}$ is invertible with $\|N^{-1}\frac{du}{dx} - I\| < 1/n$, e.g. $N = M$. Let $f = N^{-1}u - I$. Then $\frac{df}{dx} = N^{-1}\frac{du}{dx} - E$, where $E = \frac{dI}{dx}$ is the identity matrix. Since for $n > 1$, $\|N^{-1}\frac{du}{dx}(0) - I\| < 1/n$ (for $n = 1$, $\|N^{-1}\frac{du}{dx}(0) - I\| < 1/2$), we have: $\|\frac{df}{dx}(0)\| < \min\{\frac{1}{n}, \frac{1}{2}\}$. It follows that for any c satisfying $\|N^{-1}\frac{du}{dx}(0) - I\| < c < \min\{\frac{1}{2}, \frac{1}{n}\}$, there exists $a > 0$ such that: $\frac{df}{dx} \sqsupseteq h$ i.e., $f \in \int h$, where $h = [-a,a]^n \searrow [-c,c]^n$. Thus, by Proposition 4.1, if $(g_{i0})_i$ is the unique fixed point of R_f, then $I + (g_{i0})_i : \mathrm{Im}(I+f) \to [-a,a]^n$ is the local inverse of $I + f$. Hence, $(I + (g_{i0})_i) \circ N^{-1} : \mathrm{Im}(u) \to [-a,a]^n$ is the local inverse of u in $[-a,a]^n$. It follows from Proposition 4.3 that the local inverse of u is locally Lipschitz.

(ii) Suppose $u = \bigsqcup_{i\geq 0} s_i$ where $s_i = \bigsqcup_{j\in J_i} r_j \searrow [p_j, q_j]$ is a piecewise linear step function for disjoint finite indexing sets J_i ($i \geq 0$) and $\frac{du}{dx} = \bigsqcup_{i\geq 0}\beta_i$ for an increasing sequence (β_i) of linear rational step functions. We denote the midpoint matrix of β_i by M_i. As $\|M^{-1}\frac{du}{dx} - I\| < \min\{\frac{1}{2}, \frac{1}{n}\}$, we can find, by Scott continuity, the smallest $k > 0$ such that M_k is invertible and $\|M_k^{-1}\beta_k(0) - I\| < 1/n$, from which we obtain $\|M_k^{-1}\frac{du}{dx}(0) - I\| < \frac{1}{n}$. We write $N := M_k$ in accordance with item (i) above.

We obtain $N^{-1}u = \bigsqcup_{i\geq 0} N^{-1}s_i$ where $N^{-1}s_i = \bigsqcup_{j\in J_i} r_j \searrow N^{-1}[p_j, q_j]$ and

$$f = N^{-1}u - I = \bigsqcup_{i\geq 0} N^{-1}(s_i - I) = \bigsqcup_{i\geq 0} t_i,$$

where $t_i := N^{-1}s_i - I = \bigsqcup_{j\in J_i} r_j \searrow (N^{-1}[p_j, q_j] - \lambda x.x)$ is a piecewise linear step function for each $i \geq 0$. Note that, from $\mathbf{I}f = \bigsqcup_{i\geq 0}\mathcal{E}(t_i)$ we obtain:

$$\begin{aligned} R_f(\perp_0) &= -\mathbf{I}f \circ (I + \perp_0) = -\bigsqcup_{i\geq 0}\mathcal{E}(t_i) \circ (I + \perp_0) \\ &= \bigsqcup_{i\geq 0} R_{\mathcal{E}(t_i)}(\perp_0), \end{aligned}$$

and hence,

$$\begin{aligned} (g_{i0})_i &= \bigsqcup_{n\geq 0} R_f^n(\perp_0) = \bigsqcup_{n\geq 0}\bigsqcup_{i\geq 0} R^n_{\mathcal{E}(t_i)}(\perp_0) \\ &= \bigsqcup_{n\geq 0} R^n_{\mathcal{E}(t_n)}(\perp_0). \end{aligned}$$

For each piecewise linear step functions $t \in A \to \mathbf{I}B$ and $s \in B \to \mathbf{I}B$, the map $R_{\mathcal{E}(t)}(s) = -(\mathbf{I}\mathcal{E}(t)) \circ (I + s)$ is the composition of two piecewise linear step functions and is thus a piecewise linear step function. It follows by a simple induction that for each $n \geq 0$, the map $\alpha_n := R^n_{\mathcal{E}(t_n)}(\perp_0)$ is a piecewise linear step function. Finally, we have

$$u^{-1} = (I + g_0) \circ N^{-1} = \bigsqcup_{n\geq 0}(I + \alpha_n) \circ N^{-1},$$

where each $A_l := (I + \alpha_l) \circ N^{-1}$ is a piecewise linear step function, made up of single-step functions with parallelograms as their domains.

(iii) With N as above, by Proposition 4.1, $(g_{i0})_i$ is C^1 and thus from $u^{-1} = (I + (g_{i0})_i) \circ N^{-1}$ we obtain: $(u^{-1})' = (I + (g_{i0})'_i) \circ N^{-1}$, where $(g_{i0})'_i$ is the unique fixed point of $S_{(\frac{df}{dx}, (g_{i0})_i)}$. Hence, it is sufficient to show that an increasing sequence of polynomial step functions with lub $(g_{i0})'_i$ can be effectively obtained. Let α_l ($l \geq 0$) be the piecewise linear step function as in (ii) with $(g_{i0})_i = \bigsqcup_{l\geq 0}\alpha_l$. Let β_l ($l \geq 0$) be an increasing sequence of piecewise linear step functions with $u' = \bigsqcup_{l\geq 0}\beta_l$. From $f = I - N^{-1}u$ we obtain: $\frac{df}{dx} = f' = I - N^{-1}u'$. Thus, $f' = \bigsqcup_{l\geq 0}(I - N^{-1}\beta_l)$, where $\gamma_l := I - N^{-1}\beta_l$ is a piecewise linear step function for each $l \geq 0$. It follows that $\mathbf{I}f' = \bigsqcup_{l\geq 0}\mathcal{E}(\gamma_l)$, where $\mathcal{E}(\gamma_l)$ is a piecewise linear step function for each $l \geq 0$. From the Scott continuity of the functional S we get for any $h \in D^0(X_1 \to X_1)$, where $E = \frac{dI}{dx}$ is the identity matrix:

$$\begin{aligned} S_{(f', (g_{i0})_i)}(h) &= S(f', (g_{i0})_i, h) \\ &= -\mathbf{I}f' \circ (I + (g_{i0})_i) \cdot (E + h) \\ &= \bigsqcup_{l\geq 0} -\mathcal{E}(\gamma_l) \circ (I + \alpha_l) \cdot (E + h) \\ &= \bigsqcup_{l\geq 0} S(\mathcal{E}(\gamma_l), \alpha_l, h) \\ &= \bigsqcup_{l\geq 0} S_{(\gamma_l, \alpha_l)}(h). \end{aligned}$$

Hence, $S^j_{(f', (g_{i0})_i)} = \bigsqcup_{l\geq 0} S^j_{(\gamma_l, \alpha_l)}$ for any $j \geq 0$.

The uniqueness of the fixed point $S_{(f',(g_{i0})_i)}$ gives:

$$\begin{aligned}(g_{i0})'_i = g_{\min} &= \bigsqcup_{j\geq 0} S^j_{(f',(g_{i0})_i)}(\bot_1)\\ &= \bigsqcup_{j\geq 0}\bigsqcup_{l\geq 0} S^j_{(\gamma_l,\alpha_l)}(\bot_1)\\ &= \bigsqcup_{l\geq 0} S^l_{(\gamma_l,\alpha_l)}(\bot_1).\end{aligned}$$

Since, for each $l \geq 0$, α_l and γ_l are piecewise linear step functions, it follows from a simple induction that, for each $l, j \geq 0$, $S^j_{(\gamma_l,\alpha_l)}(\bot_1)$ is a polynomial step function. In particular, $B_l := S^l_{(\gamma_l,\alpha_l)}(\bot_1)$ is a polynomial step function.

(iv) Consider the two sequences, A_l and B_l converging to u^{-1} and $(u^{-1})'$ respectively as constructed in (ii) and (iii). For each, $l \geq 0$, we have $(A_l, B_l) \in D^1(\mathrm{Im}(u) \to [-a,a]^n)$ and the update [7, 8] $\mathrm{Up}(A_l, B_l)$ is a polynomial step function whose lower and upper parts are respectively the least and the greatest functions consistent with both A_l and B_l. It follows that $u^{-1} = \bigsqcup_{i\geq 0} \mathrm{Up}(A_l, B_l)$ and that the lower and upper parts give an increasing and a decreasing sequence of continuous piecewise polynomials converging in the C^1 norm to u^{-1} from below and above respectively.

7. Implicit functions

As in classical theory, the implicit function theorem in the domain-theoretic setting can be deduced from the inverse function theorem.

Theorem 7.1 Implicit Function Theorem.
Let $f : O \to \mathbb{R}^n$, where $O \subseteq \mathbb{R}^{n+k}$, be C^1 with $f(0,0) = 0$ where $(0,0) \in \mathbb{R}^n \times \mathbb{R}^k$. Assume

$$\det(\frac{\partial(f_1,\ldots,f_n)}{\partial(x_1,\ldots,x_n)})(0,0) \neq 0.$$

Then, there exists a k-dimensional open set $W \subseteq \mathbb{R}^k$ with $0 \in W$, and a unique $g : W \to \mathbb{R}^n$ with

(i) $g \in C^1$ on W, (ii) $g(0) = 0$, (iii) $f(g(t), t) = 0$ for $t \in W$.

Moreover, we have:

(iv) *Given two increasing sequences of linear rational step functions, of type $O \to \mathbf{I}\mathbb{R}^n$ and $O \to \mathbf{I}\mathbb{R}^{n\times(n+k)}$, converging to f and f' respectively, we can effectively obtain two increasing sequences of, respectively linear and polynomial, step functions converging to g and g' respectively.*

(v) *With the assumption in (vi), we can also effectively obtain an increasing sequence of polynomial step functions, whose lower and upper parts are continuous and piecewise polynomial, such that the two sequences of lower and upper parts converge in the C^1 norm, respectively from below and above, to g.*

Proof Let $u : O \to \mathbb{R}^{n+k}$ with $u_i(x,t) = f(x,t)$ for $1 \leq i \leq n$ and $u_i(x,t) = t_i$ for $n+1 \leq i \leq n+k$. Then, $u(0,0) = (0,0)$ and

$$\det u'(0,0) = \det(\frac{\partial(f_1,\ldots,f_n)}{\partial(x_1,\ldots,x_n)})(0,0) \neq 0.$$

By Theorem 6.1, there exists $a > 0$ with $[-a,a]^{n+k} \subseteq O$ such that the restriction $u : [-a,a]^{n+k} \to \mathbb{R}^{n+k}$ has a C^1 inverse $u^{-1} : \mathrm{Im}(u) \to \mathbb{R}^{n+k}$. Put $W = \pi_1(\mathrm{Im}(u))$ where $\pi_1 : \mathbb{R}^{n+k} \to \mathbb{R}^k$ is the projection to $\mathbb{R}^k$. Let $g : W \to \mathbb{R}^n$ with $g = \lambda t.u^{-1}(0,t)$. Then, as in the classical theory, $W \subseteq \mathbb{R}^k$ is open and g is the unique function that satisfies (i), (ii) and (iii). It remains to show (iv) and (v):

(iv) Since $[-a,a]^{n+k} \subseteq O$, we can restrict the domain of the step functions, given in the assumption, to $[-a,a]^{n+k}$. Suppose, therefore, we are given an increasing sequence of linear step functions $\theta_j : [-a,a]^{n+k} \to \mathbf{I}\mathbb{R}^n$, for $j \geq 0$, with $f = \bigsqcup_{j\geq 0} \theta_j$. The functional

$$s : ([-a,a]^{n+k} \to \mathbf{I}\mathbb{R}^n) \to ([-a,a]^{n+k} \to \mathbf{I}\mathbb{R}^{n+k}),$$

with $(s(h))_j = h_j$ for $1 \leq j \leq n$ and $(s(h))_j = \lambda x.x$ for $n+1 \leq j \leq n+k$ is Scott continuous and preserves linear step functions since the inclusion map $[-a,a]^{n+k} \to \mathbf{I}\mathbb{R}^{n+k}$ is clearly a linear step function. It follows that $u = \bigsqcup_{j\geq 0} s(\theta_j)$ is the lub of an increasing sequence of linear step functions.

Suppose we also have an increasing sequence of linear step functions $\psi_l : [-a,a]^{n+k} \to \mathbf{I}\mathbb{R}^{n\times(n+k)}$, for $l \geq 0$, with $f' = \bigsqcup_{l\geq 0} \psi_l$. Note that $u'_i = f'_i$ for $1 \leq i \leq n$ and $(u'_i)_j = \delta_{ij}$ for $n+1 \leq i \leq n+k$. Consider the functional $w : ([-a,a]^{n+k} \to (\mathbf{I}\mathbb{R}^{n\times(n+k)})_s) \to ([-a,a]^{n+k} \to (\mathbf{I}\mathbb{R}^{(n+k)\times(n+k)})_s)$, with $(w(\psi))_i = \psi_i$ for $1 \leq i \leq n$ and $(w(\psi))_{ij} = \delta_{ij}$ for $n+1 \leq i \leq n+k$ and $1 \leq j \leq n+k$. Then, w is Scott continuous and preserves linear step functions. It follows that we can effectively obtain an increasing sequence of piecewise linear step functions $w(\psi_l)$ with $u' = \bigsqcup_{l\geq 0} w(\psi_l)$.

By Theorem 6.1, we can effectively obtain two increasing sequence of linear step functions C_j (for $j \geq 0$) with $u^{-1} = \bigsqcup_{j\geq 0} C_j$ and D_j with $(u^{-1})' = \bigsqcup_{l\geq 0} D_l$. The functional

$$t : (\mathrm{Im}(u) \to \mathbf{I}\mathbb{R}^{n+k}) \to (W \to \mathbf{I}\mathbb{R}^n),$$

with $t(h) = \lambda t.h(0,t)$ is Scott continuous and preserves linear step functions, since partial evaluation of a linear step function, on a subset of its arguments, results in a linear step function. Thus, $g = t(u^{-1}) = \bigsqcup_{j\geq 0} t(A_j)$ is obtained effectively as the lub of an increasing sequence of step function.

For the derivative, we obtain that $(g')_i = \lambda x.(u^{-1})'_{n+i}(0,x)$ and thus effectively obtain $(g')_i = \bigsqcup_{l\geq 0} \lambda x.(D_l)_{n+i}(0,x)$, where $\lambda x.(D_l)_{n+i}(0,x)$ is a polynomial step function.

(v) As in the proof of Theorem 6.1(iv).

8 Further work

The results in the paper present a theoretical foundation for a robust CAD system. Having obtained domain-theoretic versions of the inverse and implicit function theorems, which in particular provide local C^1 approximations to an implicit surface, the next step would be to be able to patch the local pieces of an implicit surface together to obtain, in particular, a closed connected orientable manifold given by implicit equations such as $f(x,y,z) = 0$ when 0 is a regular value of f. Furthermore, the domain-theoretic framework for geometric modelling developed in [6] combined with the results in this work lead to a domain of orientable closed Lipschitz manifolds. This will synthesize the domain-theoretic framework for geometry and that for differential calculus.

References

[1] J. L. Berggren. *Episodes in the Mathematics of Medieval Islam Berggren*. Springer, 1986.

[2] J. Bloomenthal, editor. *Introduction to implicit surfaces*. Morgan Kaufmann, 1997.

[3] F. H. Clarke. *Optimization and Nonsmooth Analysis*. Wiley, 1983.

[4] B. Pavlov D. Bridges, C. Calude and D. Stefanescu. The constructive implicit function theorem and applications in physics. *Chaos, Solitons and Fractals*, 10:927–934, 1999.

[5] A. Edalat, M. Krznarić, and A. Lieutier. Domain-theoretic solution of differential equations (scalar fields). In *Proceedings of MFPS XIX*, volume 73 of *Electronuc Notes in Theoretical Computer Science*, 2003. Full paper in `www.doc.ic.ac.uk/~ae/papers/scalar.ps`.

[6] A. Edalat and A. Lieutier. Foundation of a computable solid modelling. *Theoretical Computer Science*, 284(2):319–345, 2002.

[7] A. Edalat and A. Lieutier. Domain theory and differential calculus (Functions of one variable). *Mathematical Structures in Computer Science*, 14(6):771–802, December 2004.

[8] A. Edalat, A. Lieutier, and D. Pattinson. A computational model for multi-variable differential calculus. In V. Sassone, editor, *Proc. FoSSaCS 2005*, LNCS, 2005. to appear.

[9] A. Edalat and D. Pattinson. A domain-theoretic account of picard's theorem. In *Proceedings of ICALP'04*, 2004. Full paper in `www.doc.ic.ac.uk/~ae/`.

[10] R. Howard. The inverse function theorem for Lipschitz functions (unpublished). `math.sc.edu/~howard/Notes/inverse.pdf`.

[11] F. Nevanlinna and R. Nevanlinna. *Absolute analysis*. Springer-Verlag, New York, 1973. Translated from the German by Phillip Emig, Die Grundlehren der mathematischen Wissenschaften, Band 102.

[12] J. M. Snyder. *Generative Modeling for Computer Graphics and CAD*. Academic Press, 1992.