

Interactive Computer Graphics

Lecture 17: Fire

Graphics Lecture 17: Slide 1

Amorphous Objects

Amorphous objects are still a major challenge in computer graphics.

Fire,
Smoke
Water
Clouds
&c.

They are the subject of current research - marketable in computer games and films.

Graphics Lecture 17: Slide 2

Polygons

Texture mapping polygons is fast and acceptable for short lived effects.

Overlay an flame point with a series of textured polygons (25 for a one second effect)



Graphics Lecture 17: Slide 3

Using polygons and fixing up the lighting

To make things more realistic we need to introduce secondary light sources at the point of the fire.



Graphics Lecture 17: Slide 4

Limitations of Polygons

Polygon fires are:

- Difficult to sustain for long periods
- Difficult to spread or change shape
- Difficult to introduce translucency

Though they are fast and therefore good for interactive graphics.

Graphics Lecture 17: Slide 5

Modelling Fire

Modelling fire is computationally expensive, but can produce better effects.

Flames are incandescent gases having temperature pressure and density.

The visual appearance and shape changes (due to diffusion) are functions of these.

Modelling the physics accurately is difficult

Graphics Lecture 17: Slide 6

Particle Systems

One solution is to approximate flame - and other amorphous objects - by a discrete set of small particles.

Particles can have:

- Mass
- Position
- Velocity
- Temperature
- Shape (not often used)
- Lifetime

Graphics Lecture 17: Slide 7

Particle Creation

Particles can be created according to a probability distribution.

They can be given an initial velocity and a lifetime

Depending on the simulation their movement can be determined by dynamics.

Graphics Lecture 17: Slide 8

Example - (water not fire!)



New particles created each frame
the number created per frame is normally distributed

Each particle has an initial downward velocity
again normally distributed

Each successive frame each particle is acted on by a “wind” force to the right

Graphics Lecture 17: Slide 9

Particle Dynamics

Newtonian particles: $\mathbf{f} = m \mathbf{a}$ (NB \mathbf{f} and \mathbf{a} are vectors)

$$\mathbf{a} = d\mathbf{v}/dt = d^2\mathbf{x}/dt^2$$

Given \mathbf{f} we need to find the change in position \mathbf{x} so we need to integrate.

But since we are working in frame intervals we can use a simple approximation.

$$\mathbf{v}_{t+1} = \mathbf{a}_t \Delta t + \mathbf{v}_t$$

$$\mathbf{x}_{t+1} = \mathbf{v}_t \Delta t + \mathbf{x}_t$$

Accuracy is not as important as effect!

Graphics Lecture 17: Slide 10

Practical particle system algorithm

```
for each video frame
{ generate new particles;
  remove old particles;
  for each particle
  { resolve forces by vector addition;
    calculate  $\mathbf{a}$ ,  $\mathbf{v}$ ,  $\mathbf{x}$ 
    apply rendering algorithm;
  }
}
```

Graphics Lecture 17: Slide 11

Enhancements - 1

Introduce damping:

$$\mathbf{f} = m\mathbf{a} + s\mathbf{v}$$

s is a scalar constant called damping or friction

for a gas it relates to the viscosity

An easy enhancement

Graphics Lecture 17: Slide 12

Enhancements - 2


Allow particles to collide

Problems:

Sub frame calculation required

Particle shape and size needed

Large number of particle/particle checks needed



Graphics Lecture 17: Slide 13

Simple Rendering

Project particles to the view frame, then

1. Single point for each particle
2. Blob for each particle
3. Streak for each particle
(line of motion during the frame)
4. Blend particle projection with the raster
(translucency)

Graphics Lecture 17: Slide 14

Rendering

Particle Colour

1. Make colour particle age dependent
(simple to implement)
2. Make colour temperature dependent
(requires modelling the temperature)

Graphics Lecture 17: Slide 15

The wrath of Khan

Strong contender for the worst film of 1982.

First use of a particle system to model fire:

Reeves, (1983) Particle systems - a technique for modelling a class of fuzzy objects. *Computer Graphics (SIGGRAPH 1983)* 17(3):359-376

(the next few images are from the paper)

Graphics Lecture 17: Slide 16

The Wrath of Khan - Storyboard


The genesis project

A barren planet is to be brought to life by dropping a bomb on it.

Doesn't sound very likely to me.

Graphics Lecture 17: Slide 17

The Wrath of Khan - overview




Many particle systems starting at different times to simulate the flames spreading

In total around 750,000 particles

Graphics Lecture 17: Slide 18

The Wrath of Khan - Particle Initialisation




Velocity magnitude distributed normally

Magnitude diminishes with time

Velocity direction distributed about the normal

Graphics Lecture 17: Slide 19

The Wrath of Khan - Particle Dynamics



Gravitational force pulling the particles towards the planet.

Particle lifetime short to prevent a whole parabola being visible

Graphics Lecture 17: Slide 20

The Wrath of Khan- Rendering

Planet rendered first - fake light source above the particle system to create the light emitted by the flames.

Line segment for each particle projected and drawn over the planet image, motion blur applied

Colour time dependent, yellow to red

Graphics Lecture 17: Slide 21

How did it look? Good Explosion!



Graphics Lecture 17: Slide 22

Effective flame spread



Graphics Lecture 17: Slide 23

Close up it looks like the edge of a carpet!



Graphics Lecture 17: Slide 24

Particles in Real Time - Quake

1. A pre-rendered bitmap
2. Real time rendered glow
3. Animated glowing particles



Graphics Lecture 17: Slide 25

Where Next - Modelling

Particle systems are very flexible modelling tools capable of simulating a lot of effects.

The particle modelling in the Wrath of Khan was simple and effective, but there might be some benefit in incorporating more features.

Graphics Lecture 17: Slide 26

Temperature

Gases cool as they expand - particles move slower at lower temperature.

Gas expansion is related to pressure differential.

The possibilities are endless and there have been lots of ideas tried.

Graphics Lecture 17: Slide 27

Where next - rendering

The real failure in the Wrath of Khan is the rendering.

Straight line segments are not a visual characteristic of incandescent gases


The lines are very much apparent where the particle density is low

Graphics Lecture 17: Slide 28

Bitmap Splatting

Render the particles as splats rather than lines.

Each splat has a pattern with degrees of opacity at each pixel



Laur and Hanrahan 1991

Graphics Lecture 17: Slide 29

Ray tracing - Blobs

Blobs can be thought of as 3D splats

Particles are treated as the centre of blobs and the flame dynamics can be modelled by the particles

Blobs can be warped in 3D to produce a less regular appearance for the fire.

Graphics Lecture 17: Slide 30

Ray Tracing Blobs is volume rendering


As the ray travels through a blob it adopts a proportion of the blob colour.

It may travel through the whole particle field and take up a proportion of the scene behind

Alternatively it can terminate within the particle field in places where it is denser.


Graphics Lecture 17: Slide 31

Ray tracing particle fields




Graphics Lecture 17: Slide 32

Further possibilities - Implicit Surfaces




Graphics Lecture 17: Slide 33

Further possibilities - Radiosity



Graphics Lecture 17: Slide 34

Flames with radiosity and blob shadows



Engell-Nielsen and Madsen
University of Copenhagen
(<http://www.it-e.dk/~beyond/Animations/index.html>)

Graphics Lecture 17: Slide 35