




Tutorial 7: Ray Tracing

A solid modelling system uses the following primitives:

Cylinder : C_1, C_2, r
 Sphere : C, r
 Box : A, e_1, e_2, e_3



The system is to draw the scene in orthographic projection. The viewing direction is parallel to the z axis: $(0,0,1)$.

1. Assuming a ray starts from a pixel with location (x_{pix}, y_{pix}) , devise a test for each primitive to identify simple cases when the ray cannot intersect it.
2. Use your tests to decide if the following rays:

| | (x_{pix}, y_{pix}) |
|-------|----------------------|
| Ray 1 | (32, 52) |
| Ray 2 | (32, 58) |

can be ruled out from intersecting the following objects:

| | C_1 | C_2 | r |
|------------|--------------|--------------|-----|
| Cylinder 1 | (20, 50, 50) | (50, 50, 50) | 10 |
| Cylinder 2 | (35, 55, 40) | (35, 55, 60) | 5 |

| | C | r |
|----------|--------------|-----|
| Sphere 1 | (20, 50, 50) | 10 |

| | A | e_1 | e_2 | e_3 |
|-------|--------------|------------|------------|------------|
| Box 1 | (35, 45, 40) | (15, 0, 0) | (0, 15, 0) | (0, 0, 20) |
| Box 2 | (30, 55, 40) | (5, 0, 0) | (0, -5, 0) | (0, 0, 20) |

3. For rays that intersect in Q2, what is the surface normal at the point of intersection?
4. Devise a suitable general test for use in perspective projection.