



Performance Analysis

Peter Harrison, Maria Vigliotti and Jeremy Bradley

Room 372. Email: `jb@doc.ic.ac.uk`

Department of Computing, Imperial College London

Produced with prosper and \LaTeX

Recall Jackson's theorem

- ➔ For a steady-state probability $\pi(r_1, \dots, r_N)$ of there being r_1 jobs in node 1, r_2 nodes at node 2, etc.:

$$\begin{aligned}\pi(r_1, r_2, \dots, r_N) &= \prod_{i=1}^N (1 - \rho_i) \rho_i^{r_i} \\ &= \prod_{i=1}^N \pi_i(r_i)\end{aligned}$$

where $\pi_i(r_i)$ is the steady-state probability there being n_i jobs at node i independently

PEPA and Product Form

- A product form result links the overall steady-state of a system to the product of the steady state for the components of that system
 - e.g. Jackson's theorem

- In PEPA, a simple product form can be got from:

$$P_1 \bowtie_{\emptyset} P_2 \bowtie_{\emptyset} \dots \bowtie_{\emptyset} P_n$$

- $\pi(P_1^{r_1}, P_2^{r_2}, \dots, P_n^{r_n}) = \frac{1}{G} \prod_{i=1}^n \pi(P_i^{r_i}) \dots \pi(P_n^{r_n})$

- where $\pi(P_i^{r_i})$ is steady state prob. that component P_i is in state r_i

PEPA and RCAT

- ➔ RCAT: *Reversed Compound Agent Theorem*
- ➔ RCAT can take the more general cooperation:

$$P \underset{L}{\bowtie} Q$$

- ➔ ...and find a product form, given structural conditions, in terms of the individual components P and Q

What does RCAT do?

- ➔ RCAT expresses the reversed component $\overline{P \underset{L}{\bowtie} Q}$ in terms of \overline{P} and \overline{Q} (almost)
- ➔ This is powerful since it avoids the need to expand the state space of $P \underset{L}{\bowtie} Q$
- ➔ This is useful since from the forward and reversed processes, $P \underset{L}{\bowtie} Q$ and $\overline{P \underset{L}{\bowtie} Q}$, we can find the steady state distribution $\pi(P_i, Q_i)$
- ➔ $\pi(P_i, Q_i)$ is the steady state distribution of both the forward and reversed processes (by definition)

Recall: Reversed processes

- The reversed process of a stationary Markov process $\{X_t : t \geq 0\}$ with state space S , generator matrix Q and stationary probabilities $\vec{\pi}$ is a stationary Markov process with generator matrix Q' defined by:

$$q'_{ij} = \frac{\pi_j q_{ji}}{\pi_i} \quad : i, j \in S$$

and with the same stationary probabilities $\vec{\pi}$.

Kolmogorov's Generalised Criteria

A stationary Markov process with state space S and generator matrix Q has reversed process with generator matrix Q' if and only if:

1. $q'_i = q_i$ for every state $i \in S$
2. For every finite sequence of states $i_1, i_2, \dots, i_n \in S$,

$$q_{i_1 i_2} q_{i_2 i_3} \cdots q_{i_{n-1} i_n} q_{i_n i_1} = q'_{i_1 i_n} q'_{i_n i_{n-1}} \cdots q'_{i_3 i_2} q'_{i_2 i_1}$$

where $q_i = -q_{ii} = \sum_{j : j \neq i} q_{ij}$

Finding π from the reversed process


- ➔ Once reversed process rates Q' have been found, can be used to extract $\vec{\pi}$
- ➔ In an irreducible Markov process, choose a reference state 0 arbitrarily
- ➔ Find a sequence of connected states, in either the forward or reversed process, $0, \dots, j$ (i.e. with either $q_{i,i+1} > 0$ or $q'_{i,i+1} > 0$ for $0 \leq i \leq j - 1$) for any state j and calculate:

$$\pi_j = \pi_0 \prod_{i=0}^{j-1} \frac{q_{i,i+1}}{q'_{i+1,i}} = \pi_0 \prod_{i=0}^{j-1} \frac{q'_{i,i+1}}{q_{i+1,i}}$$

Reversing a sequential component

- ➔ Reversing a sequential component, S , is straightforward:

$$\bar{S} \stackrel{\text{def}}{=} \sum_{i : R_i \xrightarrow{(a_i, \lambda_i)} S} (\bar{a}_i, \bar{\lambda}_i) . \bar{R}_i$$



Activity substitution

- ➔ We need to be able to substitute a PEPA activity $\alpha = (a, r)$ for another $\alpha' = (a', r')$:

$$(\beta.P)\{\alpha \leftarrow \alpha'\} = \begin{cases} \alpha'.(P\{\alpha \leftarrow \alpha'\}) & : \text{if } \alpha = \beta \\ \beta.(P\{\alpha \leftarrow \alpha'\}) & : \text{otherwise} \end{cases}$$

$$(P + Q)\{\alpha \leftarrow \alpha'\} = P\{\alpha \leftarrow \alpha'\} + Q\{\alpha \leftarrow \alpha'\}$$

$$(P \boxtimes_L Q)\{\alpha \leftarrow \alpha'\} = P\{\alpha \leftarrow \alpha'\} \boxtimes_{L\{\alpha \leftarrow \alpha'\}} Q\{\alpha \leftarrow \alpha'\}$$

where $L\{(a, \lambda) \leftarrow (a', \lambda')\} = (L \setminus \{a\}) \cup \{a'\}$
if $a \in L$ and L otherwise

- ➔ A set of substitutions can be applied with:

$$P\{\alpha \leftarrow \alpha', \beta \leftarrow \beta'\}$$

RCAT Conditions (Informal)

For a cooperation $P \underset{L}{\bowtie} Q$, the reversed process

$\overline{P \underset{L}{\bowtie} Q}$ can be created if:

1. Every passive action in P or Q that is involved in the cooperation $\underset{L}{\bowtie}$ must always be enabled in P or Q respectively.
2. Every reversed action \bar{a} in \overline{P} or \overline{Q} , where a is active in the original cooperation $\underset{L}{\bowtie}$, must:
 - (a) always be enabled in \overline{P} or \overline{Q} respectively
 - (b) have the same rate throughout \overline{P} or \overline{Q} respectively

RCAT Notation

In the cooperation, $P \bowtie_L Q$:

- $\mathcal{A}_P(L)$ is the set of actions in L that are also active in the component P
- $\mathcal{A}_Q(L)$ is the set of actions in L that are also active in the component Q
- $\mathcal{P}_P(L)$ is the set of actions in L that are also passive in the component P
- $\mathcal{P}_Q(L)$ is the set of actions in L that are also passive in the component Q
- \bar{L} is the reversed set of actions in L , that is
$$\bar{L} = \{\bar{a} \mid a \in L\}$$

RCAT Conditions (Formal)

For a cooperation $P \bowtie_L Q$, the reversed process

$\overline{P \bowtie_L Q}$ can be created if:

1. Every passive action type in $\mathcal{P}_P(L)$ or $\mathcal{P}_Q(L)$ is always enabled in P or Q respectively (i.e. enabled in all states of the transition graph)
2. Every reversed action of an active action type in $\mathcal{A}_P(L)$ or $\mathcal{A}_Q(L)$ is always enabled in \overline{P} or \overline{Q} respectively
3. Every occurrence of a reversed action of an active action type in $\mathcal{A}_P(L)$ or $\mathcal{A}_Q(L)$ has the same rate in \overline{P} or \overline{Q} respectively

RCAT (I)

For $P \underset{L}{\bowtie} Q$, the reversed process is:

$$\overline{P \underset{L}{\bowtie} Q} = R^* \underset{\bar{L}}{\bowtie} S^*$$

where:

$$R^* = \bar{R}\{(\bar{a}, \bar{p}_a) \leftarrow (\bar{a}, \top) \mid a \in \mathcal{A}_P(L)\}$$

$$S^* = \bar{S}\{(\bar{a}, \bar{q}_a) \leftarrow (\bar{a}, \top) \mid a \in \mathcal{A}_Q(L)\}$$

$$R = P\{(a, \top) \leftarrow (a, x_a) \mid a \in \mathcal{P}_P(L)\}$$

$$S = Q\{(a, \top) \leftarrow (a, x_a) \mid a \in \mathcal{P}_Q(L)\}$$

where the reversed rates, \bar{p}_a and \bar{q}_a , of reversed actions are solutions of Kolmogorov equations.

RCAT (II)

x_a are solutions to the linear equations:

$$x_a = \begin{cases} \bar{q}_a & : \text{if } a \in \mathcal{P}_P(L) \\ \bar{p}_a & : \text{if } a \in \mathcal{P}_Q(L) \end{cases}$$


and \bar{p}_a, \bar{q}_a are the symbolic rates of action types \bar{a} in \bar{P} and \bar{Q} respectively.

RCAT in words

To obtain $\overline{P \underset{L}{\bowtie} Q} = R^* \underset{\bar{L}}{\bowtie} S^*$:


1. substitute all the cooperating passive rates in P, Q with symbolic rates, x_{action} , to get R, S
2. reverse R and S , to get \bar{R} and \bar{S}
3. solve non-linear equations to get reversed rates, $\{\bar{r}\}$ in terms of forward rates $\{r\}$
4. solve non-linear equations to get symbolic rates $\{x_{action}\}$ in terms of forward rates
5. substitute all the cooperating active rates in \bar{R}, \bar{S} with \top to get R^*, S^*

Example: Tandem queues (I)



- ➔ Jobs arrive to node P with activity (e, γ)
- ➔ Jobs are serviced at node P with rate μ_1
- ➔ Jobs move between node P and Q with action a
- ➔ Jobs are serviced at node Q with rate μ_2
- ➔ Jobs depart Q with action d

Example: Tandem queues (II)



- ➔ PEPA description, $P_0 \bowtie_{\{a\}} Q_0$, where:

$$P_0 \stackrel{\text{def}}{=} (e, \gamma).P_1$$

$$P_n \stackrel{\text{def}}{=} (e, \gamma).P_{n+1} + (a, \mu_1).P_{n-1} \quad : n > 0$$

$$Q_0 \stackrel{\text{def}}{=} (a, \top).Q_1$$

$$Q_n \stackrel{\text{def}}{=} (a, \top).Q_{n+1} + (d, \mu_2).Q_{n-1} \quad : n > 0$$

Example: Tandem queues (III)

- ➔ Replace passive rates in cooperation with variables:

$$R = P\{(a, \top) \leftarrow (a, x_a) \mid a \in \mathcal{P}_P(L)\}$$
$$S = Q\{(a, \top) \leftarrow (a, x_a) \mid a \in \mathcal{P}_Q(L)\}$$

- ➔ Transformed PEPA model:

$$R_0 \stackrel{\text{def}}{=} (e, \gamma).R_1$$

$$R_n \stackrel{\text{def}}{=} (e, \gamma).R_{n+1} + (a, \mu_1).R_{n-1} \quad : n > 0$$

$$S_0 \stackrel{\text{def}}{=} (a, x_a).S_1$$

$$S_n \stackrel{\text{def}}{=} (a, x_a).S_{n+1} + (d, \mu_2).S_{n-1} \quad : n > 0$$

Example: Tandem queues (IV)

- ➔ Reverse components R and S to get:

$$\bar{R}_0 \stackrel{\text{def}}{=} (\bar{a}, \bar{\mu}_1) \cdot \bar{R}_1$$

$$\bar{R}_n \stackrel{\text{def}}{=} (\bar{a}, \bar{\mu}_1) \cdot \bar{R}_{n+1} + (\bar{e}, \bar{\gamma}) \cdot \bar{R}_{n-1} \quad : n > 0$$

$$\bar{S}_0 \stackrel{\text{def}}{=} (\bar{d}, \bar{\mu}_2) \cdot \bar{S}_1$$

$$\bar{S}_n \stackrel{\text{def}}{=} (\bar{d}, \bar{\mu}_2) \cdot \bar{S}_{n+1} + (\bar{a}, \bar{x}_a) \cdot \bar{S}_{n-1} \quad : n > 0$$

- ➔ Now need to find in this order:
 1. reverse rates in terms of forward rates
 2. variable x_a in terms of forward rates

Example: Tandem queues (V)

- Finding reverse rates using Kolmogorov
 - Compare forward/reverse leaving rate from states R_0, S_0 :

$$\text{exit_rate}(R_0) = \text{exit_rate}(\bar{R}_0) : \quad \bar{\mu}_1 = \gamma$$

$$\text{exit_rate}(S_0) = \text{exit_rate}(\bar{S}_0) : \quad \bar{\mu}_2 = x_a$$

- Compare rate cycles in R, \bar{R} and S, \bar{S} :

$$R_0 \rightarrow R_1 \rightarrow R_0 : \quad \gamma\mu_1 = \bar{\mu}_1\bar{\gamma}$$

$$S_0 \rightarrow S_1 \rightarrow S_0 : \quad x_a\mu_2 = \bar{\mu}_2\bar{x}_a$$

- Giving: $\bar{\gamma} = \mu_1$ and $\bar{x}_a = \mu_2$

Example: Tandem queues (VI)

- Finding symbolic rates – recall:

$$x_a = \begin{cases} \bar{q}_a & : \text{if } a \in \mathcal{P}_P(L) \\ \bar{p}_a & : \text{if } a \in \mathcal{P}_Q(L) \end{cases}$$

- In this case, $a \in \mathcal{P}_Q(L)$, so $x_a = \bar{p}_a =$ reversed rate of a -action in \bar{R}
- Thus $x_a = \bar{\mu}_1 = \gamma$
- This agrees with rate of customers leaving forward network – why?

Example: Tandem queues (VII)

→ Constructing $\overline{P \boxtimes_L Q}$

→ $\overline{P_0 \boxtimes_{\{a\}} Q_0} = R_0^* \boxtimes_{\{\bar{a}\}} S_0^*$ where:

$$R_0^* \stackrel{\text{def}}{=} (\bar{a}, \top).R_1^*$$

$$R_n^* \stackrel{\text{def}}{=} (\bar{a}, \top).R_{n+1}^* + (\bar{e}, \mu_1).R_{n-1}^* \quad : n > 0$$

$$S_0^* \stackrel{\text{def}}{=} (\bar{d}, \gamma).S_1^*$$

$$S_n^* \stackrel{\text{def}}{=} (\bar{d}, \gamma).S_{n+1}^* + (\bar{a}, \mu_2).S_{n-1}^* \quad : n > 0$$

Example: Tandem queues (VIII)

- Finding the steady state distribution:
 - Need to use the following formula:

$$\pi_j = \pi_0 \prod_{i=0}^{j-1} \frac{q_{i,i+1}}{q'_{i+1,i}}$$

...to find the steady state distribution

- First need to construct a sequence of events to a generic state (n, m) in network
 - where (n, m) represents n jobs in node P and m in node Q

Example: Tandem queues (IX)

- ➔ Generic state can be reached by:
 1. $n + m$ arrivals or e -actions to node P
(forward rate = γ , reverse rate = μ_1)
 2. followed by m departures or a -actions from node P and arrivals to node Q (forward rate = μ_1 , reverse rate = μ_2)

$$\begin{aligned}\text{Thus: } \pi(n, m) &= \pi_0 \prod_{i=0}^{n+m-1} \frac{\gamma}{\mu_1} \times \prod_{i=0}^{m-1} \frac{\mu_1}{\mu_2} \\ &= \pi_0 \left(\frac{\gamma}{\mu_1} \right)^n \left(\frac{\mu_1}{\mu_2} \right)^m\end{aligned}$$

References

- RCAT
 - *Turning back time in Markovian Process Algebra*. Peter Harrison. TCS 290(3), pp. 1947–1986. January 2003.
- Generalised RCAT: less strict structural conditions
 - *Reversed processes, product forms and a non-product form*. Peter Harrison. LAA 386, pp. 359–381. July 2004.
- MARCAT: N-way cooperation extension:
 - *Separable equilibrium state probabilities via time-reversal in Markovian process algebra*. Peter Harrison and Ting-Ting Lee. TCS, pp. 161–182. November 2005.